



SHINE-LIFT LIPSTICK 6834

Formula

A	• Diisopropyl dimer dilinoleate	10.00 %
	• Titanium dioxide	0.55 %
	• DC red 7	0.66 %
	• DC red 27	4.77 %
	• DC red 36	2.67 %
B	• Diisopropyl dimer dilinoleate	25.25 %
	• Ozokerite	12.96 %
	• Pentaerythrityl tetralaurate	10.00 %
	• Octyldoecanol	8.12 %
	• Beeswax (Cera alba)	6.82 %
	• Triisostearyl trilinoleate	5.00 %
	• Cetyl palmitate	4.50 %
	• Carnuba wax (Copernicia cerifera)	2.50 %
C	• SEPILIFT DPHP (<i>Dipalmitoyl Hydroxyproline - SEPPIC</i>)	1.00 %
	• Fragrance	5.00 %
	• Tocopheryl acetate	0.20 %

Procedure

Mill phase A using a 3-roller mill.

Melt phase B at 85°C-90°C and add, under stirring, phase A previously crushed.

Stir until complete dissolution.

Colour control is performed before introduction of phase C under stirring.

Cast into molds.

Comments

SEPILIFT DPHP

Plant-derived hydroxyproline "Lipovector". An anti-wrinkle ingredient, it moisturizes the skin and reduces the appearance of visible signs of aging (effectiveness proven in vivo). SEPILIFT DPHP firms skin tissues by stimulating the contraction of the collagen fibers. It protects dermal fibers from enzymatic lysis and exerts an anti-free radical action. SEPILIFT DPHP improves lipstick spreadability and provides comfort wearing on lips




Characteristics

Appearance
Stability

Matt stick
stable at RT/40°C

Assessment

Gloss value of 30 μ standard films (SEPPIC method 57CO042 - Reflectometer MICRO-TRIGLOSS 4520 - angle 60°) compared to control formula without SEPILIFT DPHP



Notes

Titanium dioxide : CI 77891

6834 - SEPPIC – A0201

Since this formula has not been the object of a toxicological study, the use and handling of the products proposed is purely indicative and SEPPIC accepts no responsibility for their use by another party.